arXiv:0808.0835v1 [math.OA] 6 Aug 2008

Perron-Frobenius operators and representations of the Cuntz-Krieger algebras for infinite matrices

August 6, 2008

Daniel Gonçalves[1] and Danilo Royer [2]

Abstract

In this paper we extend work of Kawamura, see [3], for Cuntz-Krieger algebras O_A for infinite matrices A. We generalize the definition of branching systems, prove their existence for any given matrix A and show how they induce some very concrete representations of O_A. We use these representations to describe the Perron-Frobenius operator, associated to an nonsingular transformation, as an infinite sum and under some hypothesis we find a matrix representation for the operator. We finish the paper with a few examples.

1 Introduction

The interactions between the theory of dynamical systems and operator algebras are one of the main venues in modern mathematics. Exploring this interplay Kawamura, see [3], recently showed that the theory of representations of the Cuntz-Krieger algebras is closely related to the theory involving the Perron-Frobenius operator. The work of Kawamura is done for the Cuntz-Krieger algebras O_A, for finite matrices A. In this paper we generalize many of the results in [3] for the Cuntz-Krieger algebras for infinite matrices (a concept introduced by Exel and Laca in [1]). For example, under some mild assumptions, we are able to give a explicit characterization of the Perron-Frobenius operator, associated to a nonsingular transformation, as a infinite sum, using a representation of an infinite Cuntz-Krieger algebra. In our efforts to generalize the notions of [3] we found two problems with the work done in there that we believe are worth mentioning. First is the necessity of an extra hypothesis in the definition of a branching function system given in [3]. The other problem is in the statement of theorem 1.2 of [3], where BA should read $A^T B$. We will deal with both these cases when introducing our generalized versions of the theory of [3].

---

[1] The author would like to thank Wael Bahsoun for bringing Kawamura´s, [3], paper to his attention.

[2] partially supported by a grant from FAPESC, Brazil

We organize the paper in the following way: In the remaining of the introduction we quickly recall the reader the main definitions of [3] and show the need for an extra hypothesis in the definition of a branching function system. In section 2, we define branching systems for infinite matrices A, which we denote by A_∞. We deal with the existence of A_∞ branching systems for any given matrix A (infinite or not) and show how they induce representations of O_A in section 3. Next, in section 4, we use the representations introduced in section 3 to describe the Perron-Frobenious operator as an infinite sum; we also present the generalizide and corrected version of theorem 1.2 of [3] in this section. We finish the paper in section 5 with a few examples.

Given a measure space (X, μ), let $L_p(X, \mu)$ be the set of all complex valued measurable functions f such that $\|f\|_p < \infty$. For a nonsingular transformation $F : X \to X$ (that is, $\mu(F^{-1}(A)) = 0$ if $\mu(A) = 0$) let $P_F : L_1(X, \mu) \to L_1(X, \mu)$ be the Perron-Frobenius operator, that is, P_F is such that

$$\int\limits_A P_F\psi(x)d\mu = \int\limits_{F^{-1}(A)} \psi(x)d\mu$$

for each measurable subset A of X, for all $\psi \in L_1(X, \mu)$. Notice that, for $\psi \in L_1(X, \mu)$, $P_F(\psi)$ is the Radon-Nikodym derivative of the measure μ_{P_F}, given by $\mu_{P_F}(A) = \int\limits_{F^{-1}(A)} \psi(x) d\mu$, with respect to μ (see [4] for more details about the Perron-Frobenius operator).

In order to describe the Perron-Frobenius operators and representations of the Cuntz-Krieger algebras, Kawamura, in [3], introduces A-branching function system on a measure space (X, μ): a family $(\{f_i\}_{i=1}^N, \{D_i\}_{i=1}^N)$ of measurable maps and measurable subsets of X respectively, together with a nonsingular transformation $F : X \to X$ such that $f_i : D_i \to f_i(D_i) = R_i$, $\mu(X \setminus \bigcup\limits_{i=1}^{N} R_i) = 0$, $\mu(R_i \cap R_j) = 0$ for all $i \neq j$, there exists the Radon-Nikodym derivative Φ_{f_i} of $\mu \circ f_i$ with respect to μ and $\Phi_{f_i} > 0$ almost everywhere in D_i for $i - 1, .., N$, $F \circ f_i = id_{D_i}$ for each $i \in \mathbb{N}$ and $\mu(D_i \setminus \bigcup\limits_{j: a_{ij}=1} R_i) = 0$, where a_{ij} are the entries of the matrix A defining O_A.

Next, a family $\{S(f_i)\}_{i=1}^N$ of partial isometries in $L_2(X, \mu)$ is defined by $S(f_i)(\phi) = \chi_{R_i} \cdot (\Phi_F)^{\frac{1}{2}} \cdot \phi \circ F$, where χ_{R_i} denotes the characteristic function of R_i, and a representation of O_A in $L_2(X, \mu)$ is obtained by defining $\pi_f(s_i) = S(f_i) \quad (i = 1..N)$, (where s_i is one of the generating partial isometry in O_A), and using the universal property of O_A. But it happens that the definition given above for an A-branching function system is not enough to guarantee that we get a representation of O_A, in fact, it is not enough to prove most of the theorems in [3]. For example, let $X = [0, 2]$, μ be the Lebesgue measure, $R_1 = [0, 1] = D_1, R_2 = [1, 2] = D_2$, $F : X \to X$ defined by $F(x) = x$ for each $x \in [0, 2]$ (so, $f_i(x) = x$ for each $x \in D_i$) and $A = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$. Following [3], $(\{f_i\}_{i=1}^2, \{D_i\}_{i=1}^2)$ is an A-branching function system, but $S(f_1)^* S(f_1)(\phi) = \chi_{[0,1]} \cdot \phi$ and $(S(f_1)S(f_1)^* + S(f_2)S(f_2)^*)(\phi) = \chi_{[0,2]} \cdot \phi$, for each $\phi \in L_2(X, \mu)$, so that $S(f_1)^* S(f_1) \neq \sum\limits_{i=1}^{2} S(f_i)S(f_i)^*$. Therefore, the existence of a representation of O_A in $L_2([0, 2], \mu)$ is not guaranteed.

As we seen, we need to add some extra hypothesis to the definition of an A-branching function system. Namely, we also have to ask that

$\mu(\bigcup_{j:a_{ij}=1} R_j \setminus D_i) = 0$, for each $i = 1, .., N$. We should mention that this extra condition is satisfied in all the examples given in [3]. With this new definition of an A-branching function system in mind, we are now able to generalize it to the countable infinite case.

2 A_∞-branching systems

For a measure space (X, μ) and for measurable subsets Y, Z of X, we write $Y \overset{\mu-a.e.}{=} Z$ if $\mu(Y \setminus Z) = 0 = \mu(Z \setminus Y)$ or equivalently, if there exists $Y', Z' \subset X$ such that $Y \cup Y' = Z \cup Z'$ with $\mu(Y') = 0 = \mu(Z')$.

Let A be an infinite matrix, with entries $A(i, j) \in \{0, 1\}$, for $(i, j) \in \mathbb{N} \times \mathbb{N}$, and let (X, μ) be a measurable space. For each pair of finite subsets U, V of $\mathbb{N}$ and $j \in \mathbb{N}$ define

$$A(U, V, j) = \prod_{u \in U} A_{uj} \prod_{v \in V} (1 - A_{vj}).$$

Definition 2.1 *An A_∞-branching system on a σ-finite measure space (X, μ) is a family $(\{f_i\}_{i=1}^{\infty}, \{D_i\}_{i=1}^{\infty})$ together with a nonsingular transformation $F : X \to X$ such that:*

1. *$f_i : D_i \to R_i$ is a measurable map, D_i, R_i are measurable subsets of X and $f_i(D_i) \overset{\mu-a.e.}{=} R_i$ for each $i \in \mathbb{N}$;*
2. *F satisfies $F \circ f_i = id_{D_i}$ $\mu - a.e.$ in D_i for each $i \in \mathbb{N}$;*
3. *$\mu(R_i \cap R_j) = 0$ for all $i \neq j$;*
4. *$\mu(R_j \cap D_i) = 0$ if $A(i, j) = 0$ and $\mu(R_j \setminus D_i) = 0$ if $A(i, j) = 1$;*
5. *for each pair U, V of finite subsets of $\mathbb{N}$ such that $A(U, V, j) = 1$ only for a finite number of j's,*

$$\bigcap_{u \in U} D_u \bigcap \bigcap_{v \in V} (X \setminus D_v) \overset{\mu-a.e}{=} \bigcup_{j \in \mathbb{N} : A(U,V,j)=1} R_j.$$

6. *There exists the Radon-Nikodym derivatives Φ_{f_i} of $\mu \circ f_i$ with respect to μ in D_i and $\Phi_{f_i^{-1}}$ of $\mu \circ f_i^{-1}$ with respect to μ in R_i.*

The existence of the Radon-Nikodym derivative Φ_{f_i} of $\mu \circ f_i$ with respect to μ in D_i together with the fact that $F \circ f_i = Id_{D_i}$ $\mu - a.e.$

imply that $f_i \circ F_{|_{R_i}} = Id_{R_i}$ $\mu - a.e.$So, the function f_i is $\mu - a.e.$ invertible, with inverse $f_i^{-1} := F_{|_{R_i}}$. These are the functions that appear in condition 6 above. If follows from the same condition that Φ_{f_i} and $\Phi_{f_i^{-1}}$ are measurable functions in D_i and R_i respectively. We will also consider these functions as measurable functions in X, defining it as being zero out of D_i and R_i, respectively.

The functions Φ_{f_i} and $\Phi_{f_i^{-1}}$ are nonnegative μ-a.e., because μ is a (positive) measure. It is possible to show (by using the following proposition) that $\Phi_{f_i} > 0$ and $\Phi_{f_i^{-1}} > 0$ $\mu-$a.e. in D_i and R_i, respectively. The same proposition shows that $\Phi_{f_i}(x)\Phi_{f_i^{-1}}(f_i(x)) = 1$ μ-almost everywhere in D_i. This equality will be used in the next section.

Proposition 2.2 *Let (X, μ) be a σ-finite measure space and Y, Z measurable subsets of (X, μ). Consider two measurable maps $f : Y \to Z$ and $g : Z \to X$, and suppose that there exists the Radon-Nikodym derivatives Φ_f of $\mu \circ f$ with respect to μ in Y and Φ_g of $\mu \circ g$ with respect to μ in Z. Suppose also that $\mu \circ f$ and $\mu \circ g$ are σ-finite. Then there exists the Radon-Nikodym derivative $\Phi_{g\circ f}$ of $\mu \circ (g \circ f)$ with respect to μ in Y and $\Phi_{g\circ f}(x) = \Phi_g(f(x))\Phi_f(x)$ $\mu - a.e$ in Y.*

Proof:

First note that $\mu \circ (g \circ f)$ is a σ-finite measure in Y.

Now, for each $E \subseteq Y$,

$$\int_E \Phi_f(x)\Phi_g(f(x))d\mu = \int_E \Phi_g(f(x))d(\mu \circ f) = \int_{f(E)} \Phi_g(x)d\mu =$$

$$= \int_{f(E)} d(\mu \circ g) = \int_E d(\mu \circ g \circ f).$$

The first and the third equality are a consequence of the Radon-Nikodym derivative. The other two follow by the change of variable theorem. So for each $E \subseteq Y$,

$$\int_E \Phi_f(x)\Phi_g(f(x))d\mu = \int_E d(\mu \circ g \circ f) = (\mu \circ g \circ f)(E).$$

So, if $\mu(E) = 0$ then $(\mu \circ g \circ f)(E) = 0$. By [2]there exists the Radon-Nikodym derivative $\Phi_{g \circ f}$ of $\mu \circ f \circ g$ with respect to μ in Y and the equality $(\mu \circ f \circ g)(E) = \int\limits_E \Phi_{g \circ f}(x) d\mu$ holds, for each $E \subseteq Y$. So, for each $E \subseteq Y$,

$$\int\limits_E \Phi_f(x)\Phi_g(f(x))d\mu = \int\limits_E \Phi_{g \circ f}(x) d\mu,$$

and therefore,$\Phi_f(x)\Phi_g(f(x)) = \Phi_{g \circ f}(x)\ \mu - a.e.$ $\square$

3 Representations of Cuntz-Krieger algebras for infinite matrices.

Representations of the Cuntz-Krieger algebras are of great importance, having applications both to operator algebras and to dynamical systems. In this section we show that for each A_∞-branching system, there exists a representation of the unital Cuntz-Krieger C*-algebra O_A on $\mathcal{B}(L_2(X, \mu))$, the bounded operators on $L_2(X, \mu)$.

Following [1], recall that the unital Cuntz-Krieger algebra of an infinite matrix A, with $A(i, j) \in \{0, 1\}$ and $(i, j) \in \mathbb{N} \times \mathbb{N}$ is the unital universal C*-algebra generated by a family $\{S_i\}_{i \in \mathbb{N}}$ of partial isometries that satisfy:

1. $S_i S_i^* S_j S_j^* = 0$ if $i \neq j$;
2. $S_i^* S_i$ and $S_j^* S_j$ commute, for all i, j;
3. $S_i^* S_i S_j S_j^* = A(i, j) S_j S_j^*$, for all i, j;
4. $\prod\limits_{u \in U} S_u S_u^* \prod\limits_{v \in V} (1 - S_v S_v^*) = \sum\limits_{j=1}^{\infty} A(U, V, j) S_j S_j^*$, for each pair of finite subsets $U, V \subseteq \mathbb{N}$ such that $A(U, V, j) := \prod\limits_{u \in U} A(u, j) \prod\limits_{v \in V} (1 - A(v, j))$ vanishes for all but a finite number of $j's$.

Theorem 3.1 *For a given A_∞-branching system (see 2.1), there exist a *-homomorphism $\pi : O_A \to \mathcal{B}(L_2(X, \mu))$ such that $\pi(S_i)\phi = \chi_{R_i} \cdot (\Phi_{f_i^{-1}})^{\frac{1}{2}} \cdot \phi \circ F$ for each $\phi \in L_2(X, \mu)$.*

Proof:

First notice that for a given $\phi \in L_2(X,\mu)$ we have that

$$\int\limits_X |\chi_{R_i}(x)\Phi_{f_i^{-1}}(x)^{\frac{1}{2}}\phi(F(x))|^2 d\mu = \int\limits_{R_i} \Phi_{f_i^{-1}}(x)|\phi(f_i^{-1}(x))|^2 d\mu =$$

$$= \int\limits_{R_i} |\phi(f_i^{-1}(x))|^2 d(\mu\circ f_i^{-1}) = \int\limits_{D_i} |\phi(x)|^2 d\mu \leq \int\limits_X |\phi(x)|^2 d\mu.$$

To obtain the second equality we have considered the Radon-Nikodym derivative of $\mu \circ f_i^{-1}$ with respect to μ in R_i and the last equality is an application of the change of variable theorem.

So, we define the operator $\pi(S_i) : \mathcal{L}(L_2(X,\mu)) \to \mathcal{L}(L_2(X,\mu))$ by

$$\pi(S_i)\phi = \chi_{R_i} \cdot (\Phi_{f_i^{-1}})^{\frac{1}{2}} \cdot (\phi \circ F),$$

for each $\phi \in L_2(X,\mu)$. By using the above computation, we see that $\pi(S_i) \in \mathcal{B}(L_2(X,\mu))$.

Our aim is to show that $\{\phi(S_i)\}_{i\in\mathbb{N}}$ satisfies the relations 1-4 which define the Cuntz-Krieger algebra O_A. With this in mind, let us first determine the operator $\psi(S_i)^*$.

For each $\phi, \psi \in L_2(X,\mu)$,

$$\langle \pi(S_i)\phi, \psi\rangle = \int\limits_X \chi_{R_i}(x)\Phi_{f_i^{-1}}(x)^{\frac{1}{2}}\phi(F(x))\overline{\psi(x)}d\mu = \int\limits_{R_i} \Phi_{f_i^{-1}}(x)^{\frac{1}{2}}\phi(f_i^{-1}(x))\overline{\psi(x)}d\mu = ...$$

...by using the change of variable theorem...

$$... = \int\limits_{D_i} \Phi_{f_i^{-1}}(f_i(x))^{\frac{1}{2}}\phi(x)\overline{\psi(f_i(x))}d(\mu\circ f_i) = ...$$

...considering the Radon derivative Φ_{f_i} of $\mu \circ f_i$...

$$... = \int\limits_{D_i} \Phi_{f_i}(x)\Phi_{f_i^{-1}}(f_i(x))^{\frac{1}{2}}\phi(x)\overline{\psi(f_i(x))}d\mu = ...$$

...by proposition 2.2...

$$... = \int\limits_{D_i} \Phi_{f_i}(x)^{\frac{1}{2}} \phi(x) \overline{\psi(f_i(x))} d\mu = \int\limits_{X} \phi(x) \Phi_{f_i}(x)^{\frac{1}{2}} \overline{\psi(f_i(x))} d\mu = \left\langle \phi, \chi_{D_i} \cdot \Phi_{f_i}^{\frac{1}{2}} \cdot (\psi \circ f_i) \right\rangle .$$

Then

$$\pi(S_i)^* \psi = \chi_{D_i} \cdot \Phi_{f_i}^{\frac{1}{2}} \cdot (\psi \circ f_i).$$

Using proposition 2.2 again, it is easy to show that

$$\pi(s_i)^* \pi(S_i) \psi = \chi_{D_i} \cdot \psi = M_{\chi_{D_i}}(\psi)$$

for each $\psi \in L_2(X, \mu)$ (that is, $\pi(S_i)^* \pi(S_i)$ is the multiplication operator by χ_{D_i}). In the same way $\pi(s_i)\pi(S_i)^* = M_{\chi_{R_i}}$.

Now we verify if $\{\pi(S_i)\}_{i \in \mathbb{N}}$ satisfies the relations 1-4, which define the C*-algebra O_A. The first relation follows from the fact that $\mu(R_i \cap R_j) = 0$ for $i \neq j$. The second one is trivial.

To see that the third relation is also satisfied, recall that if $A(i,j) = 0$ then $\mu(R_j \cap D_i) = 0$ and hence

$$\pi(S_i)^* \pi(S_i) \pi(S_j) \pi(S_j)^* = M_{\chi_{D_i}} M_{\chi_{R_j}} = M_{\chi_{D_i \cap R_j}} = 0,$$

and if $A(i,j) = 1$ then $\mu(R_j \setminus D_i) = 0$ and hence

$$\pi(S_i)^* \pi(S_i) \pi(S_j) \pi(S_j)^* = M_{\chi_{D_i}} M_{\chi_{R_j}} = M_{\chi_{D_i \cap R_j}} = M_{\chi_{R_j}} = \pi(S_j)\pi(S_j)^*.$$

So, for each $i, j \in \mathbb{N}$

$$\pi(S_i)^* \pi(S_i) \pi(S_j) \pi(S_j)^* = A(i,j) \pi(S_j) \pi(S_j)^*.$$

To verify the last relation, let U, V be finite subsets of $\mathbb{N}$ such that $A(U, V, j) = 1$ only for finitely many $j's$.

Then, by definition 2.1:5,

$$M_{\chi_{\left(\bigcap\limits_{u \in U} D_u \bigcap\limits_{v \in V} (X \setminus D_v) \right)}} = M_{\chi_{\left(\bigcup\limits_{A(U,V,j)=1} R_j \right)}} .$$

Note that

$$M_{\chi_{\left(\bigcap\limits_{u\in U} D_u \bigcap\limits_{v\in V}(X\setminus D_v)\right)}} = \prod_{u\in U} M_{\chi D_u} \prod_{v\in V}(Id - M_{\chi D_v}) =$$

$$= \prod_{u\in U} \pi(S_u)^*\pi(S_u) \prod_{v\in V}(Id - \pi(S_v)^*\pi(S_v)).$$

On the other hand,

$$M_{\chi_{\left(\bigcup\limits_{j\in\mathbb{N}:A(U,V,j)=1} R_j\right)}} = \sum_{j\in\mathbb{N}:A(U,V,j)=1} M_{\chi_{R_j}} = \sum_{j\in\mathbb{N}:A(U,V,j)=1} \pi(S_j)\pi(S_j)^*.$$

This shows that the last relation defining O_A is also verified.

So, there exist a *-homomorphism $\pi : O_A \to \mathcal{B}(L_2(X,\mu))$ satisfying $\pi(S_i)\phi = \chi_{R_i} \cdot (\Phi_{f_i^{-1}})^{\frac{1}{2}} \cdot \phi \circ F$. $\square$

The previous theorem applies only if an A_∞-branching system is given. Our next step is to guarantee the existence of A_∞-branching systems for any matrix A. First we prove a lemma, which will be helpful in some situations.

Lemma 3.2 *Let A be a infinite matrix with entries in $\mathbb{N}\times\mathbb{N}$ having no identically zero rows, (X,μ) be a measure space, and let $\{R_j\}_{j=1}^\infty$ and $\{D_j\}_{j=1}^\infty$ be families of measurable subsets of X such that*

a) $\mu(R_i \cap R_j) = 0$ *for all* $i \neq j$;

b) $X \overset{\mu-a.e.}{=} \bigcup\limits_{j=1}^{\infty} R_j$;

c) $D_i \overset{\mu-a.e.}{=} \bigcup\limits_{j\in\mathbb{N}:A(i,j)=1} R_j$;

Then conditions 4 and 5 of 2.1 are satisfied.

Proof: Condition 4 follows from *a*) and *b*). To show 5 firs we note that $X \setminus D_v \overset{\mu-a.e.}{=} \bigcup\limits_{j\in\mathbb{N}:A(v,j)=0}$. Then, given U, V finite subsets of N, we have that

$$\bigcap_{u\in U} D_u \cap \bigcap_{v\in V}(X\setminus D_v) \overset{\mu-a.e}{=} \left(\bigcup_{j\in\mathbb{N}:A(u,j)=1\forall u\in U} R_j\right) \cap \left(\bigcup_{j\in\mathbb{N}:A(v,j)=0\forall v\in V} R_j\right) \overset{\mu-a.e}{=}$$

$$\left(\bigcup_{j\in\mathbb{N}:\ \prod\limits_{u\in U} A(u,j)=1} R_j\right)\cap\left(\bigcup_{j\in\mathbb{N}:\ \prod\limits_{v\in V}(1-A(v,j))=1} R_j\right)\overset{\mu-a.e}{=}\bigcup_{j\in\mathbb{N}:A(U,V,j)=1} R_j.$$

□

Theorem 3.3 *For each infinite matrix A, without identically zero rows, there exists an A_∞-branching system in the measure space $([0,\infty),\mu)$, where μ is the Lebesgue measure.*

Proof: Consider $[0,\infty)$ with the Lebesgue measure μ. Define $R_i = [i,i+1]$ and $D_i = \bigcup\limits_{j:A(i,j)=1} R_j$. Note that $\mu(R_i\cap R_j)=0$ for $i\neq j$. Then, by the previous lemma, conditions 4 and 5 of definition 2.1 are satisfied. So, it remains to define maps $f_i : D_i \to R_i$ and $F : [0,+\infty)\to[0,+\infty)$ satisfying the conditions of definition 2.1. For a fixed $i_0\in\mathbb{N}$ we define f_{i_0} as follows. First divide the interval $\overset{\circ}{R}_{i_0}$ (where $\overset{\circ}{R}_{i_0}$ denotes the interior of R_{i_0}) in $\#\{j : A(i_0,j)=1\}$ intervals I_j. Then, define $\tilde{f_{i_0}} : \bigcup\limits_{j:A(i_0,j)=1}\overset{\circ}{R}_j \to \bigcup\limits_{j:A(i_0,j)=1}\overset{\circ}{I}_j$ such that $\tilde{f_{i_0}} : \overset{\circ}{R}_j \to \overset{\circ}{I}_j$ is a C^1-diffeomorphism. We now define $f_{i_0} : D_{i_0}\to R_{i_0}$ by

$$f_{i_0}(x)=\begin{cases}\tilde{f_{i_0}}(x) & \text{if } x\in\bigcup\limits_{j:A(i_0,j)=1}\overset{\circ}{R}_j\\[2ex] i_0 & \text{if } x\in D_{i_0}\setminus\bigcup\limits_{j:A(i_0,j)=1}\overset{\circ}{R}_j\end{cases}$$

and $F:[0,\infty)\to[0,\infty)$ by

$$F(x)=\begin{cases}\tilde{f_{i_0}}^{-1}(x) & \text{if } x\in\bigcup\limits_{j:A(i_0,j)=1}\overset{\circ}{I}_j\\[2ex] 0 & \text{if } x\in R_{i_0}\setminus\bigcup\limits_{j:A(i_0,j)=1}\overset{\circ}{I}_j\ .\end{cases}$$

Note that f_i and F are measurable maps. Moreover, $\mu\circ f_i$ and $\mu\circ f_i^{-1}$ are σ-finite measures in D_i and R_i. Next we show that there exists the Radon-Nikodym derivatives Φ_{f_i} of $\mu\circ f_i$ with respect do μ in D_i. Let $E\subseteq D_i$ be such that $\mu(E)=0$. To show that $\mu\circ f_i(E)=0$

it is enough to show that $\mu \circ f_i(E \cap (\bigcup\limits_{j:A(i,j)=1} \overset{\circ}{R_j})) = 0$, and this equality is true by [5].Then, by [2],there exist the desired nonnegative Radon-Nikodym derivative Φ_{f_i}. In the same way there exists the (nonnegative) Radon-Nikodym derivative $\Phi_{f_i^{-1}}$ of $\mu \circ f_i^{-1}$ with respect to μ in R_i. We still need to show that F is nonsingular. For this, let $A \subseteq [0, \infty)$ be such that $\mu(A) = 0$. Notice that it is enough to prove that $\mu(F^{-1}(A) \cap R_j) = 0$ for each j. Now $\mu(F^{-1}(A) \cap R_j) = \mu(f_j(A \cap D_j)) = 0$, (where the last equality follows from the fact that $\mu \circ f_j \ll \mu$ in D_j), and hence $\mu(F^{-1}(A)) = 0$ as desired.

□

Corollary 3.4 *Given an infinite matrix A, there exists a representation of O_A in $L_2([0, \infty), \mu)$ where μ is the Lebesgue measure. If A is $N \times N$ then there exists a representation of O_A in $L_2([0, N), \mu)$ where μ is the Lebesgue measure.*

4 The Perron-Frobenious Operator

We now describe the Perron-Frobenious operator using the representations introduced in the previous section.

Theorem 4.1 *Let (X, μ) be a measure space with a branching system as in definition 2.1 and let $\varphi \in L_1(X, \mu)$ be such that $\varphi(x) \geq 0$ μ-a.e..*

1. *If supp$(\varphi) \subseteq \bigcup\limits_{i=1}^{N} R_j$, then*

$$P_F(\varphi) = \sum_{i=1}^{N} \left(\pi(S_i^*)\sqrt{\varphi}\right)^2 .$$

2. *If supp$(\varphi) \subseteq \bigcup\limits_{i=1}^{\infty} R_j$, then*

$$P_F(\varphi) = \lim_{N \to \infty} \sum_{i=1}^{N} \left(\pi(S_i^*)\sqrt{\varphi}\right)^2 ,$$

where the convergence occurs in the norm of $L_1(X, \mu)$.

Proof: The first assertion follow from the fact that for each measurable set $A \subseteq X$, $\int\limits_A P_F(\varphi)(x)d\mu = \int\limits_A \sum\limits_{i=1}^{N} \left(\pi(S_i^*)\sqrt{\varphi}(x)\right)^2 d\mu$. To prove this equality, we will use the Radon-Nikodym derivative of $\mu \circ f_i$, the change of variable theorem and the fact that $F^{-1}(A) \cap R_i = f_i(A \cap D_i)$. Given $A \subseteq X$ a measurable set we have that

$$\sum_{i=1}^{N} \int\limits_A (\pi(S_i^*)\sqrt{\varphi}(x))^2 d\mu = \sum_{i=1}^{N} \int\limits_A \chi_{D_i}(x)\Phi_{f_i}(x)\varphi(f_i(x))d\mu =$$

$$= \sum_{i=1}^{N} \int\limits_{A\cap D_i} \Phi_{f_i}(x)\varphi(f_i(x))d\mu = \sum_{i=1}^{N} \int\limits_{A\cap D_i} \varphi(f_i(x))d(\mu\circ f_i) = \sum_{i=1}^{N} \int\limits_{f_i(A\cap D_i)} \varphi(x)d\mu =$$

$$= \sum_{i=1}^{N} \int\limits_{F^{-1}(A)\cap R_i} \varphi(x)d\mu = \sum_{i=1}^{N} \int\limits_{F^{-1}(A)} \chi_{R_i}\varphi(x)d\mu = \int\limits_{F^{-1}(A)} \sum_{i=1}^{N} \chi_{R_i}\varphi(x)d\mu =$$

$$= \int\limits_{F^{-1}(A)} \varphi(x)d\mu = \int\limits_A P_F(\varphi)(x)d\mu.$$

We now prove the second assertion. For each $N \in \mathbb{N}$, define $\varphi_N := \sum\limits_{i=1}^{N} \chi_{R_i} \cdot \varphi$. Note that $(\varphi_N)_{N\in\mathbb{N}}$ is an increasing sequence, bounded above by φ. Then

$$\lim_{N\to\infty} \int\limits_X P_F(\varphi_N)(x)d\mu = \lim_{N\to\infty} \int\limits_X \varphi_N(x)d\mu = ...$$

...by the Lebesgue's Dominated Convergence Theorem...

$$= \int\limits_X \varphi(X)d\mu = \int\limits_X P_F(\varphi)(x)d\mu.$$

Moreover, the sequence $(P_F(\varphi_N))_{N\in\mathbb{N}}$ is μ - a. e. increasing and bounded above by $P_F(\varphi)$.

Then,

$$\lim_{N\to\infty} \|P_F(\varphi) - P_F(\varphi_N)\|_1 = \lim_{N\to\infty} \int\limits_X |P_F(\varphi)(x) - P_F(\varphi_N)(x)|d\mu =$$

$$= \lim_{N\to\infty} \int_X P_F(\varphi)(x) - P_F(\varphi_N)(x) d\mu = 0.$$

Therefore, $\lim\limits_{N\to\infty} P_F(\varphi_N) = P_F(\varphi)$. By the first assertion, $P_F(\varphi_N) = \sum\limits_{i=1}^{N} \left(\pi(S_i^*)\sqrt{\varphi_N}\right)^2$, and a simple calculation shows that

$$\sum_{i=1}^{N} \left(\pi(S_i^*)\sqrt{\varphi_N}\right)^2 = \sum_{i=1}^{N} \left(\pi(S_i^*)\sqrt{\varphi}\right)^2.$$

So, we conclude that

$$\lim_{N\to\infty} \sum_{i=1}^{N} \left(\pi(S_i^*)\sqrt{\varphi}\right)^2 = P_F(\varphi).$$

□

Theorem 4.2 *Let A be a matrix such that each row has a finite number of 1's and let (X, μ) be an A_∞-branching system. Suppose $\mu(R_i) < \infty$ for each i (so that $\chi_{R_i} \in L_1(X, \mu)$). Moreover, suppose Φ_{f_i} is a constant positive function for each i, say $\Phi_{f_i} = b_i$ (for example, if f_i is linear). Let $W \subseteq L_1(X, \mu)$ be the vector subspace*

$$W = span\{\chi_{R_i} : i \in \mathbb{N}\},$$

that is, W is the subspace of all finite linear combinations of χ_{R_i}. Then the Perron-Frobenius operator restricted to W, $P_{F|_W} : W \to W$, has a matrix representation given by $A^T B$, where B is the diagonal infinite matrix with nonzero entries $B_{i,i} = b_i$.

Although A and B are infinite matrices, we are considering the matrix multiplication $A^T B$ as the usual multiplication for finite matrices, since B is column-finite.

Proof: Since each row z of A has a finite number of 1's, then, by definition 2.1:5, taking $Z = \{z\}$ and $Y = \emptyset$, we obtain $D_z \overset{\mu-a.e}{=} \bigcup\limits_{j:A(z,j)=1} R_j$ so that $\chi_{D_z} = \sum\limits_{j:A(z,j)=1} \chi_{R_j}$. Note that

$$P_F(\chi_{R_z}) = b_z \chi_{D_z} = \sum_{j:A(z,j)=1} b_z R_j,$$

and so the element (j, z) of the matrix representation of $P_{F|_W}$ is $b_z A(z, j)$. $\square$

5 Examples

Example 5.1 O_∞ *(O_A where all entries of the matrix A are 1).*

Consider $X = [0, 1]$ with Lebesgue measure and define $D_i = [0, 1]$, for $i = 1, 2, \ldots$. To define the R_i´s we first need to define recursively the following sequences in X: Let $a_1 = 0$, $a_i = a_{i-1} + \frac{1}{2^i}$, $i = 2, 3, \ldots$ and let $b_i = \frac{a_i + a_{i+1}}{2}$, $i = 1, 2, \ldots$. Now define $R_i = [a_{\frac{i+1}{2}}, b_{\frac{i+1}{2}}]$ for i odd and $R_i = [b_{\frac{i}{2}}, a_{\frac{i}{2}+1}]$ for i even and define a map F on X by $F(x) = \frac{x}{b_{\frac{i+1}{2}} - a_{\frac{i+1}{2}}} + \frac{a_{\frac{i+1}{2}}}{a_{\frac{i+1}{2}} - b_{\frac{i+1}{2}}}$ for $x \in R_i$, i odd and $F(x) = \frac{x}{a_{\frac{i}{2}+1} - b_{\frac{i}{2}}} + \frac{b_{\frac{i}{2}}}{b_{\frac{i}{2}} - a_{\frac{i}{2}+1}}$ for $x \in R_i$, i even. Notice that F is nothing more than an affine transformation that takes the interval R_i onto $D_i = [0, 1]$, as shown in the picture below:



Finally, let $f_i = (F_{|_{R_i}})^{-1}$. Then $(\{f_i\}_{i=1}^\infty, \{D_i\}_{i=1}^\infty)$ is an A_∞ branching system and hence induces a representation of the Cuntz-Krieger algebra O_∞.

Example 5.2

Let X be the measure space $[0,\infty)$, with the Lebesgue measure. Consider the map $F:[0,\infty)\to[0,\infty)$ defined by $F(x)=\frac{i}{2}(x-i)^2$ for $x\in[i-1,i]$ and i odd and $F(x)=[\frac{i}{2}](x-(i-1))^2$ for $x\in[i-1,i]$ and i even ($[\frac{i}{2}]$ is the least integer greater than or equal to $\frac{i}{2}$). Below we see the graph of F.



Define $R_i=[i-1,i]$ for $i=1,2,3,...$, set $D_i=[0,[\frac{i}{2}]]$ and let $f_i:D_i\to R_i$ be defined by $f_i=(F_{|_{R_i}})^{-1}$. Then $(\{f_i\}_{i=1}^{\infty},\{D_i\}_{i=1}^{\infty})$ is an A_∞ branching system. This branching system induces a representation of the C*-algebra O_A, for

$$A=\begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & 0 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & \cdots \\ 1 & 1 & 1 & 0 & \cdots \\ \vdots & \vdots & \vdots & & \end{pmatrix}.$$

References

[1] R. Exel and M. Laca, *Cuntz-Krieger algebras for infinite matrices*, J. reine. angewa. Math.**512**, (1999), 119-172.

[2] E. Hewitt and K. Stromberg, *Real and abstract analysis*, Springer-Verlag, (1975).

[3] K. Kawamura, *The Perron-Frobenius operators, invariant measures and representations of the Cuntz-Krieger algebras*, J. Math. Physics **46**, (2005).

[4] Lasota, A. and Yorke, J.A., *Exact dynamical systems and the Frobenius-Perron operator*, Trans. Am. Math. Soc., **273**, (1982), 375-384.

[5] P. J. Fernandez. *Medida e Integração*, Projeto Euclides, 2002.

D. Goncalves, Departamento de Matemática, Universidade Federal de Santa Catarina, Florianópolis, 88040-900, Brasil

Email: daemig@gmail.com

D. Royer, Departamento de Matemática, Universidade Federal de Santa Catarina, Florianópolis, 88040-900, Brasil

Email: royer@mtm.ufsc.br